



Medication Adherence: Most Important but Mostly Despised

Abdul Kader Mohiuddin *

*Alumni, Faculty of Pharmacy, Dhaka University, Bangladesh.

Corresponding author: Abdul Kader Mohiuddin, Alumni, Faculty of Pharmacy, Dhaka University, Bangladesh, Tel: +01935183385; E-mail: trymohi@yahoo.co.in

Abstract

Abbreviations: ADR: Adverse Drug Reactions; NSAIDs: Non-Steroidal Anti-Inflammatory Drugs; CDC: The Centers for Disease Control and Prevention; WHO: World Health Organization

Keywords: Patient compliance, Healthcare denials, Medication adherence, Elderly patient care, Treatment failure

Proper use of medicine or taking medicine in correct order is essential to cure any disease. According to the WHO, lack of adherence to treatment regimens leads to major problems among patients, mostly with chronic illnesses. "Right administration" depends on at least 5 right factors--right patient, right drug, right time, right dose and right route [1]. "Medicines won't work if you don't take it right"--This simple fact is not understood by most people in the world, as a result still more than half of the patients with chronic diseases in the developed world do not take their medicine correctly--says WHO [2].

Patients with chronic diseases may find it particularly difficult to adhere because they frequently need to take their medications for an extended period of time, sometimes for the remainder of their lives. There are several reasons why patients may find it challenging to adhere to treatment regimens, and CDC estimates that medication non-compliance leads to 30 to 50% of chronic disease treatment failures. Poor adherence may cause treatment outcomes to not be achieved, symptoms to worsen, and one's health to deteriorate [3].

In the UK, up to 50% of medicines are not taken as intended and 60% of NHS patients failed to receive the right treatment within 18 weeks [4-6]. Lack of medication adherence leads to poorer health outcomes, higher healthcare expenditures, increased hospitalizations, and even higher mortality rates in patients with chronic diseases [7]. Medication non-adherence alone accounts for at least 10% of hospitalizations in US, 250,000 hospitalizations in Australia and 1.1 million hospital days in France [8-10]; induces \$300 billion in annual medical costs in US, and \$125 billion in EU; causes more than 1,25,000 premature deaths in the US and 2,00,000 deaths in EU [8,11,12]. Also, two-thirds of medication-related hospital admissions in Australia are potentially preventable [9]. A recent Canadian

study found that 30% of patients stop taking their medication before it is instructed, and 25% patients do not fill their prescription or take less than prescribed [13]. Among patients with at least one preventable encounter, medication non-adherence was associated with \$679-\$898 increased preventable spending [14]. However, pharmaceutical companies around the globe lost \$637 billion in potential sales annually due to non-adherence, with \$250 billion lost for the same in the U.S. alone last year [15].

A strange similarity can be found in under-developed, developing countries and the so-called developed world in the West or the Middle-East when it comes to not taking medicine properly. More than half of the antibiotics worldwide are sold without a prescription and CDC stated, 30-50% of antibiotics prescribed in hospitals are inappropriate or unnecessary [16,17]. A recent study published by The Lancet, funded by the Bill & Melinda Gates Foundation, Wellcome Trust, states that nearly 5 million deaths worldwide in 2019 were related to bacterial resistance, which is expected to double by 2050 [18].

Globally, NSAIDs are responsible for at least 650,000 hospitalizations and 165,000 deaths and 30% of ADR related hospital admissions around the globe annually [19, 20]. Overuse of this class of drugs can cause kidney injury, and their side effects can be 3 to 4 times higher in kidney compromised patients [21]. Many studies have reported widespread misuse of these drugs in Dengue, Chikungunya

Received: November 16, 2022; **Revised:** November 22, 2022; **Accepted:** November 25, 2022

Citation: Mohiuddin AK. (2022) Medication Adherence: Most Important but Mostly Despised. *Ann Clin Case Stud Rep*, 1(1): 1-4.

Copyright: ©2022 Mohiuddin AK. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

and Covid-19 patients. Especially in Dengue or Covid-19 patients, it is more important to maintain the hydration level of the body than to bring down the fever with the pain killers. In children, the use of excess Paracetamol syrup or suppositories may cause stomach irritation, which hampers digestion and led to vomiting and ended up with hospitalization. Most hospitalizations or ICU admissions among those patients could be prevented, with few exceptions, simply by preventing dehydration at home with saline and fruit juice or simply by drinking more water [22].

More or less 40% of Covid-19 patients report sleep disturbances—use of Benzodiazepines in Covid-19 patients increases the incidence of delirium, depresses the system in patients with compromised respiratory functions, and contraindicated with some anti-viral medications [23,24]. Surprisingly, between 2020-2021, a huge increase in benzodiazepine dispensing is reported in Canada and abuse of similar drugs doubled in Italy [25]. Around 300 metric tons of morphine-type painkillers are used worldwide each year, less than 1% of which distributed to low-and-middle income countries, says the American Journal of Public Health [26]. So their misuse and related side effects are also retained by the developed world.

Health literacy and medication adherence are strongly associated. A cross-sectional study of 259 school leaders in Hong Kong carried out during the COVID-19 pandemic between April 2021 and February 2022 shows that more than 50% of participants had limited health literacy, which was strongly associated with a negative attitude towards vaccination, confusion about COVID-19-related information and secondary symptoms [27]. Earlier, a US-based study in 2020 concluded that two-thirds of the Americans will not get the COVID-19 vaccine when it is first available, while 25% report that they do not have any intention to get vaccinated at any time [28].

According to a WHO report, only half of patients in developed countries adhere to treatment guidelines for chronic diseases, which is much less in developing countries [29]. Several studies among diabetic patients in South Asian countries have shown that nearly half of patients do not adhere to their prescribed medication and are at risk of acute and long-term complications, resulting in increased hospitalization rates and medical costs [30,31].

Three-quarters of elderly patients worldwide are unable to adhere to appropriate long-term treatment regimens—due to multiple physical complications and additional medication burden [32]. Elderly patients taking at least 5 medications are at increased risk of mild cognitive impairment, dementia, falls, frailty, disability, and mortality, while ADRs are estimated to be 5% to 28% of acute geriatric medical admissions [33,34].

A recent study by the American Heart Association revealed that patients with high blood pressure do not follow

treatment guidelines because of--(1) suboptimal dosing or prescribing the wrong medication (2) lack of insurance or lack of health care access and (3) patient failure to comply prescribed medication or other lifestyle guidelines [35].

“Medical costs are barriers to adherence to proper clinical guidelines for long-term diseases in poor countries”--although discussed in many forums but forgetfulness, confusion about the duration required for medication use and mistrust about the overall efficacy of medication are among the reasons for non-adherence to diabetes management protocols in Middle Eastern countries [36].

Epilepsy is still seen in most countries of the world as an evil spirit --although two-thirds of patients can become seizure-free with adequate treatment, poor adherence to proper guidelines is a major problem for effective recovery [37].

Several identified reasons for non-adherence to treatment guidelines for chronic diseases:

- 1. Patient's socio-economic status:** low health literacy, lack of family or social support network, unstable living or homelessness, financial insecurity.
- 2. Treatment-related:** complexity and duration of treatment procedures, frequent changes in medication regimen, lack of immediate results, real or perceived unpleasant side effects, interference with lifestyle.
- 3. Health system-related:** high treatment costs, limited health system for patient education and follow-up, doctor-patient relationship, patient trust in health care, long waits, lack of patient information materials.
- 4. Patient-related:** visual-hearing and cognitive impairment, mobility and dexterity, psychological and behavioral factors, perceived risk of disease susceptibility, superstitions and stigmatization by disease, etc. [38].

Finally, it can be said that patients' knowledge and perception of the disease is the main driving force in determining their adherence to the treatment regimen. Health care providers should explore providing more effective health-education to identify patients' attitudes toward disease, trust in medications, psychological stressors, and increase medication adherence.

Financial Disclosure or Funding: N/A

Conflict of Interest: The author declares that he has no competing interests.

Informed Consent: N/A

REFERENCES

1. Grissinger M (2010) The Five Rights: A Destination Without a Map. P T 35(10): 542.

2. Brown MT, Bussell JK (2011) Medication adherence: WHO cares? *Mayo Clin Proc* 86(4): 304-314.
3. Center for Drug Evaluation and Research (2022). Why you need to take your medications as prescribed or instructed. U.S. Food and Drug Administration. Accessed on: November 05, 2022. Available online at: <https://www.fda.gov/drugs/special-features/why-you-need-to-take-your-medications-prescribed-or-instructed>
4. Barnett NL (2014) Medication adherence: where are we now? A UK perspective. *Eur J Hospital Pharm Sci Pract* 21: 181-184.
5. Denis C, Duncan P (2022) Record 6.8m People Waiting for Hospital Treatment in England. *The Guardian*. Accessed on: September 08, 2022. Available online at: www.theguardian.com/society/2022/sep/08/waiting-lists-for-routine-hospital-treatment-in-england-break-record
6. Luke A (2022) 60% of Patients Waiting 18 Wks for Treatment at Worst-performing Trust. *Mail Online*. Accessed on: March 07, 2022. Available online at: www.dailymail.co.uk/news/article-10577387/60-NHS-patients-waited-18-weeks-treatment-worst-performing-hospital-trust.html
7. Mohiuddin AK (2020) Chapter 14. Patient Compliance. *The Role of the Pharmacist in Patient Care: Achieving High Quality, Cost-Effective and Accessible Healthcare through a Team-Based, Patient-Centered Approach, Universal-Publishers* pp: 250-270. Available online at: <https://www.universal-publishers.com/book.php?method=ISBN&book=1627343083>
8. Cutler RL, Torres-Robles A, Wiecek E, Drake B, Van der Linden N, et al. (2019) Pharmacist-led medication non-adherence intervention: reducing the economic burden placed on the Australian health care system. *Patient Prefer Adherence* 13: 853-862.
9. Lim R, Ellett LMK, Semple S, Roughead E (2022) The Extent of Medication-Related Hospital Admissions in Australia: A Review from 1988 to 2021. *Drug Safety* 45(3): 249-257.
10. Wolters Kluwer (2020) Medication Nonadherence: Medicine's Weakest Link. Available online at: <https://www.wolterskluwer.com/en/expert-insights/medication-nonadherence-medicines-weakest-link>
11. Kardas P, Bago M, Barnestein-Fonseca P, Garuoliene K, Granas AG, et al. (2022) Reimbursed medication adherence enhancing interventions in 12 European countries: Current state of the art and future challenges. *Front Pharmacol* 13: 944829.
12. van Boven JF, Tsiligianni I, Potočnjak I, Mihajlović J, Dima AL, et al. (2021) European Network to Advance Best Practices and Technology on Medication Adherence: Mission Statement. *Front Pharmacol* 12: 748702.
13. Bonsu KO, Young S, Lee T, Nguyen H, Chitsike RS (2022) Adherence to Antithrombotic Therapy for Patients Attending a Multidisciplinary Thrombosis Service in Canada - A Cross-Sectional Survey. *Patient Prefer Adherence* 16: 1771-1780.
14. Zhang Y, Flory JH, Bao Y (2022) Chronic Medication Nonadherence and Potentially Preventable Healthcare Utilization and Spending Among Medicare Patients. *J Gen Intern Med* 37(14): 3645-3652.
15. Bulik BS (2016) Nonadherence Costs Pharma \$600B-plus in Annual Sales: Study. *Fierce Pharma*. Accessed on: November 22, 2016. Available online at: <https://www.fiercepharma.com/marketing/non-adherence-costs-healthcare-system-patient-outcomes-and-pharma-bottom-line>
16. Bahta M, Tesfamariam S, Weldemariam DG, Yemane H, Tesfamariam EH, et al. (2020) Dispensing of antibiotics without prescription and associated factors in drug retail outlets of Eritrea: A simulated client method. *PLoS One* 15(1): e0228013.
17. CDC (2022) Improve Antibiotic Use. Accessed on: March 09, 2022. Available online at: <https://www.cdc.gov/sixteenteen/hai/index.htm>
18. Antimicrobial Resistance Collaborators (2022) Global burden of bacterial antimicrobial resistance in 2019: A systematic analysis. *Lancet* 399(10325): 629-655.
19. Kasciuškevičiūtė S, Gumbrevičius G, Vendzelytė A, Ščiupokas A, Petrikonis K, et al. (2018) Impact of the World Health Organization Pain Treatment Guidelines and the European Medicines Agency Safety Recommendations on Non-Steroidal Anti-Inflammatory Drug Use in Lithuania: An Observational Study. *Medicina (Kaunas)* 54(2): 30.
20. Abigail D, Robson J (2016) The Dangers of NSAIDs: Look Both Ways. *Br J Gen Pract* 66(645): 172-173.
21. Lucas GNC, Leitão ACC, Alencar RL, Xavier RMF, Daher EF, et al. (2019) Pathophysiological aspects of nephropathy caused by non-steroidal anti-inflammatory drugs. *J Bras Nefrol* 41(1): 124-130.
22. Mohiuddin AK (2022) Taking Medicine in the Right Way: Most Important but Most Neglected. *Cases* 1(1): 1-3.
23. Jahrami H, BaHammam AS, Bragazzi NL, Saif Z, Faris M, et al. (2021) Sleep problems during the COVID-19

- pandemic by population: A systematic review and meta-analysis. *J Clin Sleep Med* 17(2): 299-313.
24. Ostuzzi G, Papola D, Gastaldon C, Schoretsanitis G, Bertolini F, et al. (2020) Safety of psychotropic medications in people with COVID-19: Evidence review and practical recommendations. *BMC Med* 18(1): 215.
 25. Ashish S, McMahon T, Gude J (2021) Benzodiazepine Misuse: An Epidemic within a Pandemic. *Cureus* 13(6): e15816.
 26. Bhadelia A, De Lima L, Arreola-Ornelas H, Kwete XJ, Rodriguez NM, et al. (2019) Solving the Global Crisis in Access to Pain Relief: Lessons From Country Actions. *Am J Public Health* 109(1): 58-60.
 27. Shahid R, Shoker M, Chu LM, Frehlick R, Ward H, et al. (2022) Impact of low health literacy on patients' health outcomes: A multicenter cohort study. *BMC Health Serv Res* 22(1): 1148.
 28. Alam MdM, Melhim LKB, Ahmad MT, Jemmali M (2022) Public Attitude towards COVID-19 Vaccination: Validation of COVID-Vaccination Attitude Scale (C-VAS). *J Multidiscip Healthc* 15: 941-954.
 29. Chauke GD, Nakwafila O, Chibi B, Sartorius B, Mashamba-Thompson T (2022) Factors influencing poor medication adherence amongst patients with chronic disease in low-and-middle-income countries: A systematic scoping review. *Heliyon* 8(6): e09716.
 30. Chong E, Wang H, King-Shier KM, Quan H, Rabi DM, et al. (2014) Prescribing patterns and adherence to medication among South-Asian, Chinese and white people with Type 2 diabetes mellitus: A population-based cohort study. *Diabet Med* 31(12): 1586-1593.
 31. Sohal T, Sohal P, King-Shier KM, Khan NA (2015) Barriers and Facilitators for Type-2 Diabetes Management in South Asians: A Systematic Review. *PLoS One* 10(9): e0136202.
 32. Félix IB, Henriques A (2021) Medication adherence and related determinants in older people with multimorbidity: A cross-sectional study. *Nurs Forum* 56: 834-843.
 33. Venu C, Roy K (2022) Geriatric Cognitive Decline and Polypharmacy. National Library of Medicine, StatPearls Publishing. Available online at: <https://www.ncbi.nlm.nih.gov/books/NBK574575/>
 34. Varghese D, Ishida C, Koya H (2022) Polypharmacy. National Library of Medicine, StatPearls Publishing. Available online at: <https://www.ncbi.nlm.nih.gov/books/NBK532953/>
 35. Choudhry NK, Kronish IM, Vongpatanasin W, Ferdinand KC, Pavlik VN, et al. (2022) Medication Adherence and Blood Pressure Control: A Scientific Statement from the American Heart Association. *Hypertension* 79(1): e1-e14.
 36. Alsairafi ZK, Taylor KM, Smith FJ, Alattar AT (2016) Patients' management of type 2 diabetes in Middle Eastern countries: Review of studies. *Patient Prefer Adherence* 10: 1051-1062.
 37. Lossius MI, Alfstad KÅ, Aaberg KM, Nakken KO (2017) Seponerer av Antiepileptika Ved anfallsfrihet - når og hvordan? *Tidsskrift for Den Norske Legeforening* 137(6): 451-454.
 38. Jin J, Sklar GE, Oh VMS, Li SC (2008) Factors affecting therapeutic compliance: A review from the patient's perspective. *Ther Clin Risk Manag* 4(1): 269-286.